

hep-ph/0107204

TIFR/TH/01-26

# Neutrino Anomalies in Gauge Mediated Model with Trilinear $R$ violation

Anjan S. Joshipura <sup>a</sup>, Rishikesh D. Vaidya <sup>a</sup> and Sudhir K. Vempati <sup>b</sup>

(a) *Theoretical Physics Group, Physical Research Laboratory,  
Navarangpura, Ahmedabad, 380 009, India.*

(b) *Department of Theoretical Physics, Tata Institute of Fundamental Research,  
Colaba, Mumbai 400 005, India.*

## Abstract

The structure of neutrino masses and mixing resulting from trilinear  $R$  violating interactions is studied in the presence of the gauge mediated supersymmetry breaking. Neutrino masses arise in this model at tree level through the RG-induced vacuum expectation values of the sneutrinos and also through direct contribution at 1-loop. The relative importance of these contributions is determined by the values of the strong and weak coupling constants. In case of purely  $\lambda'$  couplings, the tree contribution dominates over the 1-loop diagram. In this case, one simultaneously obtains atmospheric neutrino oscillations and quasi-vacuum oscillations of the solar neutrinos if all the  $\lambda'$  couplings are assumed to be of similar magnitudes. If  $R$  parity violation arises from the trilinear  $\lambda$  couplings, then the loop induced contribution dominates over the tree level. One cannot simultaneously explain the solar and atmospheric deficit in this case if all the  $\lambda$  couplings are of similar magnitude. This however becomes possible with hierarchical  $\lambda$  and we give a specific example of this.

## I. INTRODUCTION

Variety of observations of the solar and atmospheric neutrinos have given important information on the possible structures of neutrino masses and mixings. Based on these observations, one can infer that oscillations among three active neutrinos are likely to be responsible for the observed features of the data. These oscillations are characterized by two widely separated mass scales. At least one mixing angle involved in oscillations of atmospheric neutrinos is large [1]. The detailed data on the day-night asymmetry and recoil energy spectrum of the solar neutrinos seem to favour the presence of one more large mixing angle [2]. Thus the neutrino mass spectrum seems to be characterized by hierarchical (mass)<sup>2</sup> differences and by two large and one small mixing angles, with the small mixing angle demanded by the CHOOZ experiment [3]. Many mechanisms have been advanced to understand these features of the neutrino spectrum [4]. One of these is provided by supersymmetric theory which contains several features to make it attractive for the description of the neutrino spectrum. The lepton number violation needed to understand neutrino masses is in-built in this theory through the presence of the  $R$  parity violating couplings [5]. Moreover, it is possible to understand the hierarchical neutrino masses and large mixing among them within this framework without fine tuning of parameters or without postulating ad hoc textures for the neutrino mass matrices [6,7].

The supersymmetrized version of the standard model contains the following lepton number violating couplings:

$$W_L = \epsilon_i L_i H_2 + \lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c, \quad (1)$$

where  $L, Q, H_2$  represent the leptonic, quark and one of the Higgs doublets (up-type) respectively and  $e^c, d^c$  represent the leptonic and down quark singlets. Each of these couplings is a potential source for neutrino masses. There have been detailed studies of the effects of these couplings on neutrino masses under different assumptions [6–11]. We briefly recapitulate the relevant gross features of these studies and motivate additional work that we are going to present in this paper.

The most studied effect is that of the three bilinear mass parameters  $\epsilon_i$ , particularly in the context of the supersymmetry breaking with universal boundary conditions at a high scale [6–8,10,11]. This formalism provides a nice way of understanding suppression in neutrino mass  $m_\nu$  compared to the weak scale. The mixing among neutrinos is largely controlled in this approach by the ratios of the parameters  $\epsilon_i$ . This can be approximately described by the following matrix for a large range in parameter space of the minimal supersymmetric

standard model with radiative  $SU(2) \times U(1)$  breaking [6,7,11]:

$$U = \begin{pmatrix} c_1 & s_1 c_2 & s_1 s_2 \\ -s_1 & c_1 c_2 & c_1 s_2 \\ 0 & -s_2 & c_2 \end{pmatrix}, \quad (2)$$

where

$$s_1 = \frac{\epsilon_1}{\sqrt{\epsilon_1^2 + \epsilon_2^2}} \quad ; \quad s_2 = \sqrt{\frac{\epsilon_1^2 + \epsilon_2^2}{\epsilon_1^2 + \epsilon_2^2 + \epsilon_3^2}}. \quad (3)$$

The above matrix can nicely reproduce the small angle MSW solution together with the atmospheric neutrino anomaly. However, due to the specific structure, one cannot have two large mixing angles keeping at the same time  $|U_{e3}| \leq 0.1$  as required from the negative results of CHOOZ. Thus purely bilinear  $R$  violation with universal boundary conditions cannot account for the observed features with two large mixing angles. This may be remedied by not insisting upon the universal boundary conditions [10] or by adding right handed neutrinos [12].

In contrast to the bilinear case, the presence of trilinear interactions can allow two large angles without conflicting with the CHOOZ result. There have been several studies to determine possible set of trilinear couplings which can reproduce the observed features of neutrino masses and mixing [13]. It is not surprising that one could ‘fit’ the neutrino spectrum in these cases due to very large number of trilinear couplings. But it was realized [14,15] that gross features of the neutrino spectrum can be understood without making specific assumptions on the trilinear couplings other than requiring them to be similar in magnitude. This makes the  $R$  violation with trilinear interaction ‘predictive’ in spite of the presence of very large number of couplings.

In addition to the  $R$  violating parameters, the neutrino spectrum in these models also depends upon the nature of the supersymmetry breaking. This spectrum has been studied in the standard supergravity case with bilinear as well as trilinear couplings and in the case of gauge mediated supersymmetry breaking when  $R$  violation is only through the bilinear terms in eq.(1). The supersymmetry breaking generically introduces two different types of contributions to neutrino masses. The presence of terms linear in the sneutrino field in the scalar potential induces a vacuum expectation value ( $v_{ev}$ ) for the former which mix neutrinos with neutralinos and lead to neutrino masses. In addition to this ‘tree level’ contribution, the trilinear terms in the superpotential also lead to neutrino masses at the 1-loop level.

Both the types of contributions are present in models with purely bilinear or purely trilinear terms in the superpotential at a high scale. In case of the trilinear couplings, the running of the couplings to lower scale generates in the scalar potential couplings linear in the sneutrino  $vev$  and lead to a tree level contribution which is often neglected in the literature.

The relative importance of the tree level and loop induced contributions to neutrino masses in case of trilinear interactions was studied in [15,16] in the context of the standard supergravity (mSUGRA) models with universal boundary conditions at a high scale. It was concluded that the tree level contribution dominates over the loop for large ranges in the parameters of the model. This results [15] in the following hierarchy in neutrino masses if all the trilinear  $\lambda'$  couplings are assumed to be similar in magnitudes:

$$\frac{m_{\nu_2}}{m_{\nu_3}} \approx \frac{m_{\text{loop}}}{m_0 + m_{\text{loop}}} \frac{m_s}{m_b} . \quad (4)$$

The parameters  $m_{\text{loop}}$  and  $m_0$  characterize the strength of the 1-loop and the RG-induced tree level contributions respectively and are determined by soft SUSY breaking parameters.  $m_s$  and  $m_b$  denote the strange quark and the bottom quark masses respectively. It is possible to obtain the vacuum or the MSW solution (large angle) to solar neutrino problem in this context by choosing the SUSY parameters in appropriate range [15].

An attractive alternative to the standard supergravity induced SUSY breaking is provided by the gauge mediated SUSY breaking [21]. Neutrino mass spectrum has been studied in gauge mediated models with trilinear R-violation by Choi *et al.* in [22]. Their study has been confined to non-minimal models of this category. The minimal model in this category called the Minimal Messenger Model(MMM) [23] has only two free parameters and is more predictive than the standard SUGRA based models and the models studies in [22]. The two free parameters of the model determine all the soft terms at the high scale  $\sim$  a few hundred TeV, where SUSY breaking occurs. Thus this model implies very constrained spectrum for neutrino masses. This constrained spectrum has been shown [7] to be inadequate for simultaneous solution of the solar and atmospheric neutrino anomalies in case of purely bilinear  $R$  violation. In this work, we wish to study neutrino masses in the minimal messenger model in the presence of purely trilinear  $R$  violation. This would mean that both the scale at which SUSY breaking occurs as well as the boundary conditions at the high scale would be sufficiently different from the mSUGRA scenario which has been studied in [15]. We have studied the neutrino mass spectrum in the MMM for two separate cases, namely purely  $\lambda'$  couplings and purely  $\lambda$  couplings. Such a choice has been made for simplicity. The  $\lambda'$  couplings with comparable magnitude are argued to describe neutrino spectrum well. In contrast, we find that if all the  $\lambda$  couplings are of similar strength, then one cannot describe

the neutrino spectrum well and one needs to postulate somewhat inverse hierarchy among them. We give a specific example with hierarchical  $\lambda$  which reproduces the observed features of the neutrino spectrum.

Within the Minimal Messenger Model, the soft supersymmetry breaking terms are decided by the gauge quantum numbers of the fields. As we will demonstrate later, this significantly alters the hierarchy within the neutrino mass states. In particular, we find that the  $m_0$  dominates over  $m_{\text{loop}}$  in case of the  $\lambda'$  couplings but the situation is reversed when the  $R$  violation occurs through  $\lambda$  couplings. This feature is characteristic of the gauge mediated scenario and is quite distinct from all the earlier studies.

We discuss the basic formalism in the next section which also contains analysis of the effect of the trilinear  $\lambda'$  couplings. The third section has detailed study of the  $\lambda$  couplings and we end with a discussion in the last section.

## II. FORMALISM

We consider the following trilinear interaction in this section:

$$W_{\mathcal{R}_p} = \lambda'_{ijk} L_i Q_j d_k^c, \quad (5)$$

where  $i, j, k$  are generation indices. In spite of very large number of these couplings, one could determine the neutrino masses and mixing in terms of small number of parameters if one assumes that all the trilinear couplings are similar in magnitude. The basic formalism was developed in [15] and we recapitulate here the relevant parts.

The neutrinos obtain their masses from two different contributions in this case. The  $\lambda'$  couplings generate radiative masses through exchange of the down squarks at the 1-loop level. In addition to this, the trilinear interactions also radiatively generate soft SUSY breaking terms which are linear in the sneutrino fields. These terms lead to additional contribution to neutrino masses which can dominate over the first contribution. The second contribution follows from the RG improved scalar potential [15,16]:

$$V_{\text{scalar}} = m_{\tilde{\nu}_i}^2 |\tilde{\nu}_i|^2 + m_{H_1}^2 |H_1^0|^2 + m_{H_2}^2 |H_2^0|^2 + \left[ m_{\nu_i H_1}^2 \tilde{\nu}_i^* H_1^0 - \mu B_\mu H_1^0 H_2^0 - B_{\epsilon_i} \tilde{\nu}_i H_2^0 + H.c \right] + \frac{1}{8} (g_1^2 + g_2^2) (|H_1^0|^2 - |H_2^0|^2)^2 + \dots, \quad (6)$$

where we have used the standard notation for the SM fields and their masses, with  $B_{\epsilon_i}$  and  $m_{\nu_i H_1}^2$  representing the bilinear lepton number violating soft terms. Minimization of the above potential leads to the sneutrino *vevs*:

$$\langle \tilde{\nu}_i \rangle = \frac{B_{\epsilon_i} v_2 - m_{\nu_i H_1}^2 v_1}{m_{L_i}^2 + \frac{1}{2} M_Z^2 \cos 2\beta} \quad , \quad (7)$$

where  $v_1$  and  $v_2$  stand for the *vevs* of the Higgs fields  $H_1^0$  and  $H_2^0$  respectively <sup>1</sup>.

These *vevs* vanish at a high scale since we are assuming only trilinear  $L$  violating interactions. They however get generated at the weak scale. The magnitudes of the parameters  $B_{\epsilon_i}$  and  $m_{\nu_i H_1}^2$  and hence the sneutrino *vevs* at the weak scale are determined by solving the renormalization group (RG) equations satisfied by them. These RG equations are presented in Appendix I. The general solution of these equations can be parameterized as

$$\begin{aligned} B_{\epsilon_i} &= \lambda'_{ipp} h_p^D \kappa_{ip} \quad , \\ m_{\nu_i H_1}^2 &= \lambda'_{ipp} h_p^D \kappa'_{ip} \quad , \end{aligned} \quad (8)$$

where  $\kappa, \kappa'$  are dependent on the soft terms appearing in the RHS of the respective RG equations and  $h^D$  are down type quark yukawa.

The sneutrino *vevs* break  $R$  parity and lead to mixing of neutrinos with neutralinos. This in turn leads to neutrino masses. For small sneutrino *vevs*, the neutrino mass matrix follows from the seesaw approximation and is given by [20]:

$$\mathcal{M}_{ij}^0 = \frac{\mu(cg^2 + g'^2) \langle \tilde{\nu}_i \rangle \langle \tilde{\nu}_j \rangle}{2(-c\mu M_2 + 2 M_W^2 c_\beta s_\beta (c + \tan \theta_W^2))} \quad , \quad (9)$$

where  $c = 5g'^2/3g^2 \sim 0.5$ ,  $M_W$  is the  $W$  boson mass,  $\mu$  is the mass term for Higgs/Higgsino,  $M_2$  is mass term for a gaugino, and the Weinberg angle is represented by  $\theta_W$ . Assuming generation independence of the terms  $\kappa, \kappa'$  which was found to be a very good approximation in [15], we can rewrite the above mass matrix as

$$\mathcal{M}_{ij}^0 \equiv m_0 \lambda'_{ipp} h_p^D \lambda'_{jmm} h_m^D \quad , \quad (10)$$

---

<sup>1</sup>These sneutrino *vevs* are derived from the tree level scalar potential. Corrections from the one-loop effective potential can significantly shift these naive tree level values [17]. For neutrino phenomenology these corrections would be important in regions in the parameter space where two contributions to the sneutrino *vev* cancel each other [18]. Such regions are not encountered in MMM parameter space in which we are interested. Moreover, we are approximately including the effect of 1-loop corrections by dynamically choosing soft parameters at appropriate scale in the manner discussed in Refs. [17,19].

where the parameters  $p$  and  $m$  are summed over the three generations and  $m_0$  now contains the dependence of the tree level mass on the soft SUSY breaking parameters. Only one neutrino attains mass through this mechanism. The other neutrinos attain mass at the 1-loop level. The complete 1-loop structure of the neutrino masses has been discussed in [18]. In the present case, the most dominant contributions are from diagrams having  $\lambda'$  couplings at both the vertices. The mass matrix generated by these diagrams is given by

$$\mathcal{M}_{ij}^l = \frac{3}{16\pi^2} \lambda'_{ilk} \lambda'_{jkl} v_1 h_k^D \sin \phi_l \cos \phi_l \ln \frac{M_{2l}^2}{M_{1l}^2} . \quad (11)$$

In the above,  $\sin \phi_l \cos \phi_l$  determines the mixing of the squark-antisquark pairs and  $M_{2l}^2$  and  $M_{1l}^2$  represent the eigenvalues of the standard  $2 \times 2$  mass matrix of the down squark system. The indices  $l$  and  $k$  are summed over. As the mixing  $\sin \phi_l \cos \phi_l$  is proportional to  $h_l^D$ , we rewrite the 1-loop contribution as,

$$\mathcal{M}_{ij}^l = m_{\text{loop}} \lambda'_{ilk} \lambda'_{jkl} h_k^D h_l^D , \quad (12)$$

where  $m_{\text{loop}}$  is independent of the  $R$  violation and solely depends on the MSSM parameters.

The total neutrino mass matrix is given by,

$$\mathcal{M}^\nu = \mathcal{M}^0 + \mathcal{M}^l \quad (13)$$

which can be rewritten in the following form when  $\mathcal{O}(h_1^{D^2}, h_2^{D^2})$  terms are neglected:

$$\mathcal{M}_{ij}^\nu \approx (m_0 + m_{\text{loop}}) a_i a_j + m_{\text{loop}} h_2^D h_3^D A_{ij} \quad (14)$$

where  $a_i = \lambda'_{ipp} h_p^D$  ( $p$  summed over generations) and

$$A_{ij} = \lambda'_{i23} \lambda'_{j32} + \lambda'_{i32} \lambda'_{j23} - \lambda'_{i22} \lambda'_{j33} - \lambda'_{i33} \lambda'_{j22} . \quad (15)$$

To derive the eigenvalues of the total matrix  $\mathcal{M}^\nu$ , we recognise that a) The first matrix on the RHS of eq.(14) has only one non-zero eigenvalue ; b) The dominant terms in the total matrix  $\mathcal{M}^\nu$  of  $\mathcal{O}(h_3^{D^2})$  are present only in the first matrix . Moreover, as we will show below  $m_{\text{loop}} \ll m_0$  in the Minimal Messenger Model in the purely  $\lambda'$  case. Hence approximate eigenvalues can be derived up to  $\mathcal{O}(\frac{m_s m_{\text{loop}}}{m_b(m_0 + m_{\text{loop}})})$ , neglecting the high order corrections. The detailed derivation of the eigenvalues and the mixing matrix has been presented in [15]. These eigenvalues are given as,

$$\begin{aligned} m_{\nu_1} &\sim m_{\text{loop}} h_2^D h_3^D \delta_1 \\ m_{\nu_2} &\sim m_{\text{loop}} h_2^D h_3^D \delta_2 \\ m_{\nu_3} &\sim (m_0 + m_{\text{loop}}) \sum_i^3 a_i^2 \\ &\sim (m_0 + m_{\text{loop}}) h_3^{D^2} \sum_i^3 \lambda'_{i33}{}^2 \end{aligned} \quad (16)$$

where

$$\begin{aligned}\delta_1 &= (c_1^2 A'_{11} - 2c_1 s_1 A'_{12} + s_1^2 A'_{22}) , \\ \delta_2 &= (s_1^2 A'_{11} + 2c_1 s_1 A'_{12} + c_1^2 A'_{22}) .\end{aligned}\tag{17}$$

The entries  $A'_{ij}$  are the elements of the matrix  $A' = U_{\lambda'}^T A U_{\lambda'}$  where

$$U_{\lambda'} = \begin{pmatrix} c_2 & s_2 c_3 & s_2 s_3 \\ -s_2 & c_2 c_3 & c_2 s_3 \\ 0 & -s_3 & c_3 \end{pmatrix} ,\tag{18}$$

with

$$s_2 = \frac{a_1}{\sqrt{a_1^2 + a_2^2}} \quad ; \quad s_3 = \sqrt{\frac{a_1^2 + a_2^2}{a_1^2 + a_2^2 + a_3^2}} .\tag{19}$$

The total mixing is given as [15],

$$\begin{aligned}K' &= U_{\lambda'} U_{\lambda'}' \\ &= \begin{pmatrix} c_1 c_2 - s_1 s_2 c_3 & s_1 c_2 + c_1 s_2 c_3 & s_2 s_3 \\ -s_2 c_1 - s_1 c_2 c_3 & -s_1 s_2 + c_1 c_2 c_3 & c_2 s_3 \\ s_1 s_3 & -s_3 c_1 & c_3 \end{pmatrix} ,\end{aligned}\tag{20}$$

where the 1-2 mixing angle  $\theta_1$  is given by,

$$\tan 2\theta_1 = \frac{2A'_{12}}{A'_{22} - A'_{11}} .\tag{21}$$

From eq.(15), we see that in the limit of exact degeneracy of the  $\lambda'$  couplings, the parameters  $A_{ij}$  would be zero. In this case, only one neutrino becomes massive in spite of the inclusion of the loop corrections. The 1-2 mixing also remains undetermined in this case. There is no reason *a priori* for the exact equality of  $\lambda'$  and non-zero but similar value for these parameters determine the 1-2 mixing to be large (see eq.(21)) and also generates mass for the other two neutrinos.

### A. MMM and neutrino anomalies

The parameters  $m_{\text{loop}}$  and  $m_0$  appearing in eq.(16) are independent of the details of the  $R$  violation and get determined by the soft SUSY breaking terms. We assume throughout



that SUSY breaking is mediated by the standard gauge interactions [21]. We work in the so-called minimal messenger model [23]. It is characterized by a messenger sector with a pair of superfields which transform vector-like under a gauge group chosen to be  $SU(5)$  for unification purposes. SUSY breaking is characterized by a singlet chiral superfield whose scalar and the auxiliary components acquire *vevs* breaking supersymmetry. This breaking is communicated to the visible sector by loop diagrams. The gauginos acquire masses at the 1-loop level which are given as,

$$M_i(X) = \tilde{\alpha}_i(X) \Lambda g(x) , \quad (22)$$

where  $X = \lambda < S >$  is the supersymmetric mass of the scalar and fermionic components of the singlet superfield,  $\Lambda$  is the ratio  $\frac{F_S}{\langle S \rangle}$ ,  $F_S$  being the *vev* of the auxiliary field of the singlet. The parameter  $x$  is defined as  $\frac{\Lambda}{X}$ . The scalars acquire masses at the two loop level. They are given by

$$m_i^2 = 2\Lambda^2 \left( C_3^i \tilde{\alpha}_3^2(X) + C_2^i \tilde{\alpha}_2^2(X) + \frac{3}{5} Y_i^2 \tilde{\alpha}_1^2(X) \right) f(x) . \quad (23)$$

where  $C_3, C_2$  are the quadratic casimirs of the gauge groups  $SU(3)$  and  $SU(2)$  respectively and  $Y_i$  being the hypercharge of the scalar field  $i$ , with  $i = \{Q_j, D_j, U_j, L_j, E_j, H_1, H_2\}$ , where  $j = 1, 2, 3$  is the generation index. The functions  $f(x), g(x)$  are given in [24] and the dependence of the soft masses on these functions is minimal. In the present analysis, we follow [19] and choose  $x = \frac{1}{2}$ . Since the dependence of the soft masses on  $x$  is minimal, a different choice of  $x$  would not significantly modify the results presented here.

The major feature characterizing the model is the absence of  $A$ -terms and the  $B$  terms in the soft potential at the scale  $X$ .

$$A(X) = 0 , \quad B(X) = 0 . \quad (24)$$

Thus, the entire soft spectrum of this model gets essentially determined by one parameter  $\Lambda$ . The parameters  $\tan\beta$  and  $\mu$  are fixed at the weak scale by requiring the breaking of the  $SU(2) \times U(1)$  symmetry. The relevant equations following from the tree level potential are given by

$$\begin{aligned} \sin 2\beta &= \frac{2B_\mu \mu}{m_{H_1}^2 + m_{H_2}^2 + 2\mu^2} \\ \mu^2 &= \frac{m_{H_1}^2 - m_{H_2}^2 \tan^2 \beta}{\tan^2 \beta - 1} - \frac{1}{2} M_Z^2 \end{aligned} \quad (25)$$

where all the parameters on the RHS of the above equations are evolved to the weak scale using the MSSM RGE. Because of the boundary condition eq.(24), the value of the  $B$

parameter at the weak scale remains small. This pushes  $\tan\beta$  to very large values in this model [19,25]. The above equations are strictly valid in case of the MSSM and  $R$  violating trilinear couplings can give corrections to these. The smallness of neutrino masses however require very tiny trilinear couplings of  $O(10^{-4} - 10^{-5})$  which would contribute negligible to eqs.(25). We thus continue to use eqs.(25).

The neutrino masses in eq.(16) are strongly hierarchical in the limit  $m_{\text{loop}} \ll m_0$ . Specifically, one obtains from eq.(16),

$$\frac{m_{\nu_2}}{m_{\nu_3}} \sim \frac{m_{\text{loop}}}{m_0} \frac{m_s}{m_b} \left( \frac{\delta_2}{\sum_i (\lambda'_{i33})^2} \right) \quad (26)$$

where  $m_s$  and  $m_b$  represent the strange and the bottom quark masses. For all the  $\lambda'$  of similar magnitudes, this ratio is completely determined by the parameter  $\Lambda$ .

We can determine the above ratio by exactly solving the RG equations, (65) . Before doing this, it is instructive to study the approximate expressions obtained when one neglects the  $Q^2$  dependence of the parameters appearing on the RHS of the RGE. In other words, we neglect the effect of running of the soft masses (from high scale,  $X$  to weak scale  $M_Z$ ) appearing in the expressions of the RGE as well as for neutrino masses. Instead, we take them to be their high scale values given by eqs.(23). Noting that more dominant contribution to sneutrino  $vev$  comes from the  $m_{\nu_i H_1}^2$  term in eq.(7) and integrating the corresponding RGE for  $m_{\nu_i H_1}^2$  in the above approximation, one finds in this simplifying case :

$$m_o \sim \left( \frac{2 \cos \beta}{3\pi^2} \right)^2 \frac{M_W^2}{\Lambda} \frac{\tilde{\alpha}_3^4(X)}{\tilde{\alpha}_2^5(X)} \left( \ln \frac{X^2}{M_Z^2} \right)^2 \quad (27)$$

The  $m_{\text{loop}}$  defined in eq.(11) has the following approximate form in the same approximation as above where we neglect the running of the soft masses.

$$m_{\text{loop}} \sim \left( \frac{v^2 \mu}{\Lambda^2} \right) \frac{\cos \beta \sin \beta}{8 \pi^2 \tilde{\alpha}_2^2(X)} \quad (28)$$

Eq.(27) clearly demonstrates that the often neglected [13,14] RG induced contribution dominates over the loop contribution in the present case of MMM just as in the case of the supergravity induced breaking [15,26]. One would have naively thought that this will not be the case in gauge mediated model since the running of masses in this case (signified by  $t \equiv \ln(\frac{X^2}{M_Z^2}) \sim 11$  in eq.(27)) is over much smaller range than in the supergravity case where  $t \equiv \ln(\frac{M_{GUT}^2}{M_Z^2}) \sim 66$ . But smallness of  $t$  in MMM is compensated by the largeness of the ratio  $\frac{m_Q^2}{m_L^2}$  (signified by the factor)  $\frac{\tilde{\alpha}_3^4}{\tilde{\alpha}_2^4}$ . As a result of which the value of  $m_0$  here can be comparable to the corresponding value [15] in supergravity case. From the expressions we

see that dependence on the  $\Lambda$  is more severe for the 1-loop mass,  $m_{\text{loop}}$  compared to the tree level contribution,  $m_0$ . However, the  $\mu$  parameter in the numerator increases approximately linearly with  $\Lambda$  [19]. This makes the  $\Lambda$  dependence of the both the contributions essentially the same. The ratio  $\frac{m_{\text{loop}}}{m_0}$  following from eqs.(27,28) is given as:

$$\frac{m_{\text{loop}}}{m_0} \sim \left(\frac{\pi^2}{3}\right) \left(\frac{\mu \tan \beta v^2}{M_W^2 \Lambda t^2}\right) \left(\frac{\tilde{\alpha}_2^3(X)}{\tilde{\alpha}_3^4(X)}\right) \quad (29)$$

From the above we see that the dependence of the ratio on  $\Lambda$  is essentially determined by the way the  $\mu$  parameter scales with respect to  $\Lambda$ . This leads to a very mild dependence of the ratio on  $\Lambda$ . Such  $\Lambda$  ‘independence’ has also been seen in the case of bilinear R-violating models in MMM [7]. For  $\Lambda = 100$  TeV,  $t = 2 \ln\left(\frac{X^2}{M_Z^2}\right) = 10.6$ ,  $\tan \beta = 46$  and  $\mu = 400$  GeV, from the above we see that the ratio is 0.39. From eqs.(27,28) we see that the typical order of magnitude for the ratio of the mass eigenvalues, eq.(26) is :

$$\frac{m_{\nu_2}}{m_{\nu_3}} \sim 10^{-3} \quad (30)$$

However the above expressions are approximate. We have determined this ratio exactly by solving the relevant RG equations numerically. For this, we closely follow the work of [19] where two loop RGE for the  $B$  parameter were used for fixing the sign of  $\mu$  parameter at the weak scale. As mentioned above, we have chosen  $x = \frac{1}{2}$ . This fixes  $X = 2\Lambda$ . Following [19], the decoupling scale ‘ $Q_0$ ’ is defined to be the geometrical mean of  $m_Q^2(X)$ ,  $m_U^2(X)$ . Such a choice of  $Q_0$  would be approximately equivalent to adding corrections due to the complete 1-loop effective potential. The gauge couplings and the Yukawa couplings are evolved from  $M_Z$  to  $Q_0$  using the Standard Model (SM) beta-functions and using MSSM beta-functions from  $Q_0$  to the high scale,  $X$ . All the soft masses and soft parameters defined at  $X$  are evolved from  $X$  to the decoupling scale  $Q_0$ . The ratio  $\frac{m_{\nu_2}}{m_{\nu_3}}$ , determined following the above procedure is plotted in Figure 1 for  $\Lambda$  varying from (50 – 150) TeV.

From the figure we see that the ratio of the eigenvalues,

$$\frac{m_{\nu_2}}{m_{\nu_3}} \approx (1 - 2) \times 10^{-4} \left(\frac{\delta_2}{\sum_i (\lambda'_{i33})^2}\right) \quad (31)$$

is typically around the expected value, eq.(30). While the ratio shown in Fig. 1 is completely fixed by the value of  $\Lambda$ , the neutrino mass ratio is uncertain by a number of  $\mathcal{O}(1)$  which is related to the trilinear parameters.

We now turn to discussing feasibility of the model for the simultaneous description of the solar and atmospheric data. The two generation analysis of each of these experiments constrain the value of the relevant  $(mass)^2$  difference and mixing angle. The allowed values for the parameters [28–30] are displayed below:

| Anomaly    | Solution         | $\Delta m^2(eV^2)$                     | $\tan^2 \theta$           |
|------------|------------------|----------------------------------------|---------------------------|
| Solar      | MSW-SMA          | $(2 - 10) \times 10^{-6}$              | $(1 - 20) \times 10^{-4}$ |
|            | MSW-LMA          | $(2 - 80) \times 10^{-5}$              | 0.2 - 4.                  |
|            | LOW-QVO          | $1 \times 10^{-10} - 4 \times 10^{-7}$ | 0.1 -8.                   |
|            | Vacuum (Just-So) | $(4 - 12) \times 10^{-12}$             | 0.1 -7.                   |
| Anomaly    |                  | $\Delta m^2(eV^2)$                     | $\sin^2 2\theta$          |
| Atmosphere |                  | $(1 - 8) \times 10^{-3}$               | 0.83 - 1.                 |

At present, global analysis including the recent results from the day/night recoil electron energy spectrum and charged current rates from SNO of the solar neutrinos from super-Kamiokande [2] favours the MSW-LMA solution but the other solutions are also allowed [28,30]. It follows from eq.(31) and the table that the most natural solution for the solar neutrino problem is through the vacuum oscillations but quasi-vacuum solution can also be obtained, if the  $\lambda'$  dependent factor in eq.(31) is somewhat large (e.g.  $\sim 5$ ) instead of exactly being one. The most preferred LMA solution cannot however be obtained.

The mixing among neutrinos is essentially controlled by ratios of trilinear couplings. This mixing is given by eq.(20). It is seen from this equation that a choice of angles  $s_{1,2,3}$  is possible which reproduces two large and one small mixing angles as required by the present data. As an example, consider the choice

$$c_1 = c_3 = s_1 = s_3 = \frac{1}{\sqrt{2}} ; s_2 \sim .13 . \quad (32)$$

This gives

$$\begin{aligned} \sin^2 2\theta_A &\equiv 4K_{\mu 3}'^2(1 - K_{\mu 3}'^2) \approx 0.99 , \\ \sin^2 2\theta_S &\equiv 4K_{e1}'^2 K_{e2}'^2 \approx 0.95 , \\ K_{e3}' &= 0.09 . \end{aligned} \quad (33)$$

This choice reproduces the required mixing angles and also satisfies the CHOOZ constraint.

### III. MODELS WITH $\lambda_{IJK}$

In this section, we discuss the structure of neutrino masses and mixing in the presence of only trilinear  $\lambda$  couplings. The lepton number violating part of the superpotential is given as,

$$W_{\mathcal{L}} = \lambda_{ijk} L_i L_j e_k^c . \quad (34)$$

There are two basic changes here compared to the last section. Firstly, the  $\lambda_{ijk}$  are antisymmetric in the first two indices restricting their total number to nine. This strongly restricts neutrino mass structure and one does not get phenomenologically consistent spectrum when all the trilinear couplings are assumed to be similar in magnitudes. Secondly, unlike in the  $\lambda'$  case, the loop induced contribution dominates over the tree level SUSY breaking in the minimal messenger model. Such dominant loop contribution has been earlier seen in some particular regions of the parameter space in the mSUGRA framework [15,22]. Here this dominance follows generically.

We present here a formalism which is similar to the spirit of the previous section in order to understand the basic features of the neutrino mass matrix. As before, the RG improved effective soft potential contains terms  $B_{\epsilon_i}$  and  $m_{\nu_i H_1}^2$  which break the lepton number. These terms are generated due to the presence of  $\lambda$  couplings in the superpotential. At the weak scale, the magnitude of these depends on the respective RGE, which we have presented in Appendix I. The solutions of eqs.(66) can be written as,

$$B_{\epsilon_i} = \lambda_{ipp} h_{pp}^E \tilde{\kappa}_{ip}, \quad (35)$$

$$m_{\nu_i H_1}^2 = \lambda_{ipp} h_{pp}^E \tilde{\kappa}'_{ip}, \quad (36)$$

where ( $i \neq j$ ) due to the anti-symmetric nature of the  $\lambda$  couplings and  $\tilde{\kappa}$  and  $\tilde{\kappa}'$  represent the dependence on the soft masses in the RGE and  $h^E$  are the charged lepton yukawa. Following similar arguments as in Section II, the presence of these terms in the scalar potential would lead to a tree level neutrino mass matrix of the following form:

$$\mathcal{M}_{ij}^0 = m_0 b_i b_j, \quad (37)$$

where  $m_0$  contains the dependence on the soft terms and  $b_i$  are given as,  $b_i = \lambda_{ipp} h_{pp}^E$  ( $i \neq p$ ). In addition to the tree level mass, the presence of  $\lambda$  couplings also gives rise to contributions at the 1-loop level. Assuming only canonical 1-loop contributions to be the most dominant contributions the 1-loop level mass matrix has the form:

$$\mathcal{M}_{ij}^l = \frac{1}{16\pi^2} \lambda_{ilk} \lambda_{jkl} v_1 h_k^E \sin \phi_l \cos \phi_l \ln \frac{M_{2l}^2}{M_{1l}^2}, \quad (38)$$

where  $\sin \phi_l \cos \phi_l$  and  $M_{1l}, M_{2l}$  represent the mixing and the eigenvalues respectively of the standard  $2 \times 2$  stau slepton mass matrix. Following the previous section, we rewrite the above as,

$$\mathcal{M}_{ij}^l = m_{\text{loop}} \lambda_{ilk} \lambda_{jkl} h_l^E h_k^E \quad (39)$$

In writing the above, we have implicitly assumed the anti-symmetric nature of the couplings. The total neutrino mass matrix is given as,

$$\mathcal{M} = \mathcal{M}^0 + \mathcal{M}^l \quad (40)$$

The above can be rewritten in the form:

$$\mathcal{M}_{ij} = (m_0 + m_{\text{loop}}) b_i b_j + m_{\text{loop}} h_3^E h_2^E B_{ij} + \mathcal{O}(h_2^{E^2}, h_2^E h_1^E), \quad (41)$$

where we have neglected  $\mathcal{O}(h_2^{E^2}, h_2^E h_1^E)$  contributions to the mass matrix. The matrix  $B$  is given as

$$B = \begin{pmatrix} \lambda_{132} \lambda_{123} - \lambda_{133} \lambda_{122} & \lambda_{123} \lambda_{232} - \lambda_{122} \lambda_{233} & \lambda_{132} \lambda_{323} - \lambda_{133} \lambda_{322} \\ \lambda_{123} \lambda_{232} - \lambda_{233} \lambda_{122} & 0 & 0 \\ \lambda_{132} \lambda_{323} - \lambda_{133} \lambda_{322} & 0 & 0 \end{pmatrix} \quad (42)$$

We diagonalise the total matrix  $\mathcal{M}$  in the same manner as for the  $\lambda'$  case and as described in [15]. However we do not make any assumption on the relative magnitude of  $m_0$  and  $m_{\text{loop}}$ . The approximate eigenvalues correct up to  $\mathcal{O}(h_2^{E^2})$  are derived as,

$$\begin{aligned} m_{\nu_1} &\approx m_{\text{loop}} h_3^E h_2^E \eta_1 + \mathcal{O}(h_2^{E^2}) \\ m_{\nu_2} &\approx m_{\text{loop}} h_3^E h_2^E \eta_2 + \mathcal{O}(h_2^{E^2}) \\ m_{\nu_3} &\approx (m_0 + m_{\text{loop}}) \sum_i^3 b_i^2 \\ &\approx (m_0 + m_{\text{loop}}) h_3^{E^2} \eta_3 + \mathcal{O}(h_3^E h_2^E) \end{aligned} \quad (43)$$

The parameters  $\eta_1, \eta_2, \eta_3$  are given by,

$$\begin{aligned} \eta_1 &= (c_1^2 B'_{11} - 2c_1 s_1 B'_{12} + s_1^2 B'_{22}) \\ \eta_2 &= (s_1^2 B'_{11} + 2c_1 s_1 B'_{12} + c_1^2 B'_{22}) \\ \eta_3 &= \sum_{i=1,2} \lambda_{i33}^2 \end{aligned} \quad (44)$$

The parameters  $B'_{ij}$  are elements of the matrix,  $B' = U_\lambda^T B U_\lambda$  where,  $U_\lambda$  has the same form as  $U_{\lambda'}$  in eq.(18) with the angles now given by,

$$s_2 = \frac{b_1}{\sqrt{b_1^2 + b_2^2}} \quad , \quad s_3 = \sqrt{\frac{b_1^2 + b_2^2}{b_1^2 + b_2^2 + b_3^2}} \quad (45)$$

The total mixing matrix is given as in eq.(20) with the 1 – 2 mixing angle given by,

$$\tan 2\theta_1 = \frac{2B'_{12}}{B'_{22} - B'_{11}} \quad (46)$$

In the above, we have made no assumption on the relative magnitudes of  $m_0$  and  $m_{\text{loop}}$ . Only assumption is almost equality of all the  $\lambda$  couplings. Most definitions are formally the same as in case of the  $\lambda'$  couplings but the physical consequences are quite different. This follows from the expressions of mixing angles  $s_{2,3}$  in eq.(45). If no hierarchy is assumed among the  $\lambda$  couplings then due to antisymmetry of  $\lambda$  we get,

$$\frac{b_2}{b_3} \approx \frac{b_1}{b_3} \approx \mathcal{O}\left(\frac{h_2^E{}^2}{h_3^E{}^2}\right)$$

As a result, eq.(45) implies  $c_3 \sim \mathcal{O}(\frac{m_\mu}{m_\tau})$  and the mixing matrix, eq.(20) can be written as

$$\begin{aligned} K &= U_\lambda U'_\lambda \\ &= \begin{pmatrix} c_1 c_2 & s_1 c_2 & s_2 \\ -s_2 c_1 & -s_1 s_2 & c_2 \\ s_1 & c_1 & 0 \end{pmatrix} + \mathcal{O}\left(\frac{m_\mu}{m_\tau}\right). \end{aligned} \quad (47)$$

The equations for  $\eta_1, \eta_2$  and  $\tan 2\theta_1$  take the following approximate forms in this case:

$$\begin{aligned} \eta_1 &\approx s_1^2 c_2^2 B_{11} + 2B_{12} c_2 s_1 (c_1 - s_1 s_2) \\ \eta_2 &\approx c_1^2 c_2^2 B_{11} - 2B_{12} c_1 s_2 (s_1 + c_1 s_2) \end{aligned} \quad (48)$$

$$\tan 2\theta_1 \approx \frac{-2B_{12}}{B_{11} c_2 - 2s_2 B_{12}} \quad (49)$$

With the hierarchical masses, the effective mixing angles  $\theta_A$  and  $\theta_{\text{CHOOZ}}$  probed by the atmospheric data and the CHOOZ experiment respectively are given by

$$\sin^2 2\theta_A = 4K_{\mu 3}^2 (1 - K_{\mu 3}^2) \quad \sin^2 2\theta_{\text{CHOOZ}} \approx 4K_{e 3}^2 (1 - K_{e 3}^2) .$$

Eq.(47) implies that these two mixing angles are equal in conflict with the observation.

The above derivation has not assumed any specific mechanism for the supersymmetry breaking and thus the conclusions are valid in supergravity scenario as well as in the MMM case considered here. It is quite interesting that in spite of the presence of nine independent  $\lambda$  parameters, one cannot explain solar and atmospheric neutrino anomaly as long as these parameters are of similar magnitudes. Departure from equalities of  $\lambda$  can lead to explanations of these anomalies and we shall present a specific example in the next section.

### A. MMM and Neutrino Anomalies

In this sub-section, we first determine the numerical values of the parameters  $m_{\text{loop}}$  and  $m_0$  which enter the neutrino mass. The dependence on trilinear couplings is factored out in defining these parameters. But their numerical values are quite different here compared to the  $\lambda'$  case studied in the earlier section. This follows since the soft sfermion masses which determine these parameters are quite different in these two cases ( see RG equations in the appendix ).

In models with gauge mediated SUSY breaking, the soft masses are proportional to the gauge quantum numbers they carry. Thus particles with strong interactions have much larger soft masses compared to the weakly interacting particles, as is evident from the eqs.(22,23). The effect of the gauge couplings also trickles down to the parameters  $B_{\epsilon_i}, m_{\nu_i H_1}^2$  through the corresponding soft masses present in their respective renormalization group equations. In the presence of purely  $\lambda'$  interactions, the strong coupling determines the magnitudes of  $B_{\epsilon_i}, m_{\nu_i H_1}^2$  at the weak scale and in turn the tree level mass as is evident from eq.(27). The loop contribution is still however determined by the weak coupling, eq.(28). It is thus the interplay between the strong coupling and the weak coupling which lead to a suppressed loop mass in the case of purely  $\lambda'$  couplings. In the case of pure  $\lambda$  couplings, the squark masses do not enter the definition of  $m_0$  and  $m_{\text{loop}}$  and both of these are determined by the weak coupling. However the dependence on the power of the weak coupling is different. Following the same method as described in the  $\lambda'$  case (above eq.(27)), an estimate of the parameter  $m_0$  is given by,

$$m_0 = \left( \frac{\cos \beta}{8\pi^2} \right)^2 \frac{M_W^2}{\Lambda} \frac{1}{\tilde{\alpha}_2(X)} \left( \ln \frac{X^2}{M_Z^2} \right)^2 \quad (50)$$

The 1-loop contribution  $m_{\text{loop}}$  can also be estimated in the similar manner as,

$$m_{\text{loop}} = \left( \frac{v^2 \mu}{\Lambda^2} \right) \frac{\cos \beta \sin \beta}{24 \pi^2 \tilde{\alpha}_2^2(X)} \quad (51)$$



The ratio  $\frac{m_{\text{loop}}}{m_0}$  is then given by,

$$\frac{m_{\text{loop}}}{m_0} \approx \left( \frac{8\pi^2}{3} \right) \left( \frac{v^2 \mu \tan \beta}{t^2 \Lambda M_W^2} \right) \frac{1}{\tilde{\alpha}_2(X)} \quad (52)$$

where  $t = \ln \left( \frac{X^2}{M_Z^2} \right)$ . The above is typically of  $\mathcal{O}(10^2)$  for  $\Lambda = 100$  TeV,  $\mu = 400$  GeV,  $\tan \beta = 46$  which shows that the tree level mass is much suppressed compared to the 1-loop mass. Comparing the above equation with that of the corresponding one for the  $\lambda'$  case, eq.(29), we see that the absence of strong-weak interplay in this case leads to a much larger ratio.

In a general mSUGRA inspired scenario, the tree level mass is much larger compared to the 1-loop mass [15] for large range in MSSM parameters, irrespective of the nature of  $R$  parity breaking. In the present case, the relative importance of loop and the tree level contributions is sensitive to the nature of  $R$  violation as demonstrated above. This feature arises not as a consequence of running of soft masses but due to difference in the relevant RG equations in case of  $\lambda$  and  $\lambda'$  couplings and the boundary conditions themselves which strongly depend on the gauge couplings in these models.

We have determined the ratio  $\frac{m_{\text{loop}}}{m_0}$  by solving the relevant RG equations numerically in the manner described in section (3). This ratio is plotted versus  $\Lambda$  in Fig.2 for  $\Lambda = (50 - 150)$  TeV. In this range,

$$\frac{m_{\text{loop}}}{m_0} = 25 - 45. \quad (53)$$

as expected from eq.(52).

The dominance of  $m_{\text{loop}}$  has the following important implication. The neutrino mass ratio following from eq.(43) is given by

$$\frac{m_{\nu_2}}{m_{\nu_3}} = \frac{m_{\text{loop}}}{m_0 + m_{\text{loop}}} \frac{m_\mu \eta_2}{m_\tau \eta_3} \quad (54)$$

In the previous case, the hierarchy in  $m_{\text{loop}}$  and  $m_0$  resulted in strong hierarchy between neutrino masses. As a result, one could only obtain vacuum or quasi-vacuum solution for the solar neutrino. Here due to  $m_{\text{loop}} \gg m_0$ , hierarchy in neutrino mass is much weaker, and we have,

$$\frac{m_{\nu_2}}{m_{\nu_3}} \approx \frac{m_\mu}{m_\tau} \sim 5 \times 10^{-2} \quad (55)$$

We can thus easily get the scale relevant for the LMA solution of the solar neutrino. As already argued the mixing pattern is not appropriate if all the  $\lambda$  couplings are similar. This is no longer true if  $\lambda$  obey some specific hierarchy as we discuss now.

## B. Illustrative Model

We assume that the couplings  $\lambda_{123}, \lambda_{233}, \lambda_{322}$  dominate over the rest and neglect the latter. Moreover we assume that non-zero couplings satisfy the following hierarchy,

$$\frac{\lambda_{233}}{\lambda_{322}} \approx \mathcal{O}\left(\frac{h_2^E}{h_3^E}\right) \quad ; \quad \frac{\lambda_{123}}{\lambda_{322}} \leq \mathcal{O}\left(\frac{h_2^E}{h_3^E}\right) \quad (56)$$

We do not have strong theoretical reasons to assume the above hierarchy. The following considerations should therefore be viewed as an example which leads to the successful explanation of the neutrino anomalies.

The tree level mass matrix in the presence of these couplings is given by,

$$\mathcal{M}_s^0 = m_0 \begin{pmatrix} 0 & 0 & 0 \\ 0 & \lambda_{233}^2 h_3^{E\ 2} & \lambda_{233} h_3^E \lambda_{322} h_2^E \\ 0 & \lambda_{233} h_3^E \lambda_{322} h_2^E & \lambda_{322}^2 h_2^{E\ 2} \end{pmatrix} \quad (57)$$

The 1-loop level mass matrix is given by,

$$\mathcal{M}_s^l = m_{\text{loop}} \begin{pmatrix} 0 & \lambda_{123} h_2^E \lambda_{232} h_3^E & 0 \\ \lambda_{123} h_2^E \lambda_{232} h_3^E & \lambda_{233}^2 h_3^{E\ 2} & \lambda_{232} h_2^E \lambda_{323} h_3^E \\ 0 & \lambda_{232} h_2^E \lambda_{323} h_3^E & \lambda_{322}^2 h_2^{E\ 2} \end{pmatrix} . \quad (58)$$

In view of the hierarchy in eq.(56), the total mass matrix has the following simple form

$$\mathcal{M}_s \approx \begin{pmatrix} 0 & x & 0 \\ x & A & A \\ 0 & A & A \end{pmatrix} , \quad (59)$$

where

$$\begin{aligned} A &\equiv (m_0 + m_{\text{loop}}) \lambda_{233}^2 h_3^{E\ 2} , \\ x &\equiv -m_{\text{loop}} \lambda_{322} \lambda_{123} h_2^E h_3^E . \end{aligned} \quad (60)$$

We shall assume  $x \leq A$  which is consistent with the hierarchy in eq.(56). One can diagonalise the above matrix:

$$R_{12}(\theta_{12}) \ R_{13}(\theta_{13}) \ R_{23}(\pi/4) \ \mathcal{M}_s \ [R_{12}(\theta_{12}) \ R_{13}(\theta_{13}) \ R_{23}(\pi/4)]^T \approx \text{Diag.}(m_{\nu_1}, m_{\nu_2}, m_{\nu_3}) .$$

Here,  $R_{ij}$  denotes rotation in the  $ij^{th}$  plane with angle  $\theta_{ij}$ . We have neglected a small contribution of  $\mathcal{O}(\frac{x}{2\sqrt{2}A})$  to the 2 – 3 mixing angle in the above derivation. The mixing angles are given by

$$\begin{aligned}\tan 2\theta_{12} &\approx \frac{4\sqrt{2}A}{x} , \\ \tan 2\theta_{13} &\approx \frac{x}{A\sqrt{2}} .\end{aligned}\tag{61}$$

The eigenvalues can be approximately written as

$$\begin{aligned}m_{\nu_1} &\approx \frac{x}{\sqrt{2}} - \frac{x^2}{8A} , \\ m_{\nu_2} &\approx -\frac{x}{\sqrt{2}} - \frac{x^2}{8A} , \\ m_{\nu_3} &\approx 2A + \frac{x^2}{2A} .\end{aligned}\tag{62}$$

It is seen from the last two equations that all the mixing angles and the masses are predicted in terms of only two parameters namely,  $x$  and  $A$ . The atmospheric mixing angle is predicted to be around  $\pi/4$  and the other two mixing angles can be expressed in terms of the solar and atmospheric scales. Using eqs.(61,62) we find,

$$\begin{aligned}\tan 2\theta_{solar} &\approx \frac{4\sqrt{2}A}{x} \approx 4\sqrt{2} \left( \frac{8\sqrt{2}\Delta_{solar}}{\Delta_A} \right)^{-1/3} , \\ \tan 2\theta_{CHOOZ} &\approx \frac{x}{A\sqrt{2}} \approx \frac{1}{\sqrt{2}} \left( \frac{8\sqrt{2}\Delta_{solar}}{\Delta_A} \right)^{1/3} .\end{aligned}\tag{63}$$

Choosing  $\Delta_A \sim 3 \cdot 10^{-3} \text{ eV}^2$  and  $\Delta_{solar} \sim (2 - 50) \cdot 10^{-6} \text{ eV}^2$  we get,

$$\begin{aligned}\frac{x}{A} &\approx (0.19 - 0.57) , \\ \tan^2 \theta_{solar} &\approx (0.93 - 0.81) , \\ U_{e3}(CHOOZ) &\approx (0.06 - 0.19) .\end{aligned}\tag{64}$$

The predictions for the mixing angles are in very good agreement with the observations which prefer large mixing angle solution for the solar neutrino. The required value of  $\frac{x}{A}$  is also consistent with the assumed hierarchy in eq.(56) among the trilinear couplings.

#### IV. DISCUSSION

We have discussed the structure of neutrino masses and mixing in the Minimal messenger model (MMM) of gauge mediated supersymmetric breaking with purely trilinear R

violating interactions. We considered two specific cases of purely  $\lambda'$  interactions and purely  $\lambda$  interactions for simplicity. The model contains very large number of parameters even under this simplifying assumptions. Remarkably, it is possible to make meaningful statement on the neutrino spectrum in spite of the presence of many unknown parameters if all these parameters are assumed similar in magnitude. This is a natural assumption in the absence of any specific symmetry to restrict the trilinear  $R$  parity violation. It is not always easy to justify this specific choice, e.g use of a  $U(1)$  symmetry which uses Froggatt -Nielsen [FN] mechanism to obtain quark and lepton masses tend to forbid all the trilinear terms altogether [33].

In the case where only  $\lambda'$  couplings are present, one naturally gets large mixing between the neutrino states. Further, the MMM offers a very constrained structure giving rise to a large hierarchy between the masses  $\sim O(10^{-2})$  for all the parameter space. The model is suitable for obtaining simultaneously solutions for atmospheric neutrino problem and quasi-vacuum oscillations.

Assumption of approximate equality of  $\lambda$  couplings in case with only  $\lambda$  couplings, leads to very constrained and phenomenologically inconsistent pattern for neutrino mixing. This conclusion follows on general grounds and it is true even if SUSY breaking is induced by supergravity interactions. It is quite interesting that one can arrive at this strong conclusions in spite of the presence of many unknown parameters by simply assuming them to be of similar magnitude.

One can obtain consistent picture of neutrino anomalies if  $\lambda$  couplings are assumed to be hierarchical. We provided an example which leads to two large and one small mixing and correct hierarchy between the solar and atmospheric neutrino scales.

One interesting result of this analysis is the interplay between the sneutrino  $vev$  induced contribution and the loop induced contribution to neutrino masses. In the context of supergravity induced SUSY breaking, it has been shown that the large logarithmic factors induced due to RG scaling enhance the sneutrino  $vev$  induced contribution compared to the loop contribution. We showed that this remains true even in the gauge mediated models of SUSY breaking in case of the trilinear  $\lambda'$  couplings. In the mSUGRA model, the dominance of tree level mass follows simply from the large factor  $t = \ln \frac{M_{GUT}^2}{M_Z^2}$  in sneutrino  $vev$  generated by running of soft parameters. In the present case, the tree level dominance occurs essentially due to boundary conditions. In case of  $\lambda'$  couplings,  $m_0$  is determined by squark masses which depend upon  $\alpha_3(X)$ .  $m_0$  dominates over loop contribution in this case. For  $\lambda$  couplings,  $m_0$  is determined by sleptons rather than by squarks masses. Due to their

dependence on weak couplings, slepton masses are much smaller than squark masses. As a consequence,  $m_0$  is suppressed compared to  $\lambda'$  case. This results in loop dominance if R is violated by  $\lambda$  couplings.

The neutrino mass hierarchy strongly depends on the ratio  $\frac{m_{\text{loop}}}{m_0}$ . In case of the tree level dominance (purely  $\lambda'$  couplings) one obtains strong hierarchy and vacuum solution while the case with loop mass dominating corresponds to milder hierarchy and the LMA MSW solution. This analysis along with other similar analysis [15,16,22] therefore underlines the need of including both the contributions to neutrino masses in a proper way.

**Acknowledgments:** We warmly acknowledge discussions with U. Chattopadhyay which have been very helpful.

## Appendix I

Here we present the Renormalization Group Equations (RGE) for the soft parameters  $B_{\epsilon_i}$  and  $m_{\nu_i H_1}^2$  for the two cases considered in this work: either purely  $\lambda'$  couplings or purely  $\lambda$  couplings are the sources of lepton number violation in the superpotential. These equations have been derived using the general formulae given in [31]. These equations can also be found in [16], whereas the equations for the standard soft parameters appearing in the RHS of the equations can be found in many papers such as [32]. In writing the below eqs.(65)(eqs.(66)), we have neglected  $\mathcal{O}(\lambda'^2)$  ( $\mathcal{O}(\lambda^2)$ ) corrections. The notation is as described in the text.

*$\lambda'$  couplings in the superpotential:*

$$\begin{aligned} \frac{dB_{\epsilon_i}(t)}{dt} &= B_{\epsilon_i}(t) \left( -\frac{3}{2}Y_t(t) - \frac{1}{2}Y_i^E(t) + \frac{3}{2}\tilde{\alpha}_2(t) + \frac{3}{10}\tilde{\alpha}_1(t) \right) \\ &\quad - \frac{3}{16\pi^2}\mu(t)\lambda'_{ijj}(t)h_{jj}^d(t) \left( \frac{1}{2}B_\mu(t) + A_{ijj}'(t) \right) \\ \frac{dm_{\nu_i H_1}^2(t)}{dt} &= m_{\nu_i H_1}^2(t) \left( -\frac{1}{2}Y_i^E(t) - \frac{3}{2}Y_b(t) - \frac{1}{2}Y_\tau(t) \right) - \frac{3}{32\pi^2}\lambda'_{ipp}(t)h_{pp}^d(t) \left( m_{H_1}^2(t) \right. \\ &\quad \left. + m_{L_i}^2(t) + 2 m_{Q_p}^2(t) + 2 A_{ipp}'(t)A_{pp}^D(t) + 2 m_{D_p}^2(t) \right) \end{aligned} \quad (65)$$

*$\lambda$  couplings in the superpotential:*

$$\begin{aligned} \frac{dB_{\epsilon_i}(t)}{dt} &= B_{\epsilon_i}(t) \left( -\frac{3}{2}Y_t(t) - \frac{1}{2}Y_i^E(t) + \frac{3}{2}\tilde{\alpha}_2(t) + \frac{1}{2}\tilde{\alpha}_1(t) \right) \\ &\quad - \frac{1}{16\pi^2}\mu(t)\lambda_{idd}(t)h_{dd}^E(t) \left( A_{idd}^\lambda(t) + \frac{1}{2}B_\mu(t) \right) \\ \frac{dm_{\nu_i H_1}^2(t)}{dt} &= m_{\nu_i H_1}^2(t) \left( -\frac{1}{2}Y_i^E(t) - \frac{1}{2}Y_\tau(t) - \frac{3}{2}Y_b(t) \right) - \frac{1}{32\pi^2}\lambda_{ijj}(t)h_{jj}^E(t) \left( m_{H_1}^2(t) \right. \\ &\quad \left. + m_{L_i}^2(t) + 2 m_{L_j}^2(t) + 2 A_{ijj}^\lambda(t)A_{jj}^E(t) + 2 m_{E_j}^2(t) \right) \end{aligned} \quad (66)$$

- [1] S. Fukuda *et al.* [Super-Kamiokande Collaboration], *Phys. Rev. Lett.* **85** (2000) 3999
- [2] S. Fukuda *et al.* [Super-Kamiokande Collaboration], hep-ex/0103033.
- [3] M. Apollonio *et al.* [CHOOZ Collaboration], *Phys. Rev. D* **61** (2000) 012001 [hep-ex/9906011].
- [4] For reviews of various models, please see, A. Yu. Smirnov, hep-ph/9901208; R. N. Mohapatra, hep-ph/9910365; A. S. Joshipura, *Pramana*, **54** (2000) 119.
- [5] L. J. Hall and M. Suzuki, *Nucl. Phys.* **B 231** (1984) 419.
- [6] K. S. Babu and A. S. Joshipura, Talks at the Trieste Conf. on ‘Quarks and Leptons: Masses and Mixing’, Oct.1996, Oct,1997.
- [7] A. S. Joshipura and S. K. Vempati, *Phys. Rev.* **D60** (1999) 095009.
- [8] T.Banks *et. al.*, *Phys. Rev. D* **52** (1995) 5319; R. Hempfling, *Nucl. Phys.* **B478** (1996) 3; H. Nilles and N. Polonsky *Nucl. Phys.* **B484** (1997) 33; M. A. Diaz, J. C. Romao and J. W. F. Valle, *Nucl. Phys.* **B524** (1998) 23. E. J. Chun, S. K. Kang, C. W. Kim, *Nucl. Phys.* **B544** (1999) 89; E. J. Chun and J. S. Lee, *Phys. Rev. D* **60** (1999) 075006; E. J. Chun, *Phys. Lett.* **B505**(2001) 155.
- [9] A. Yu. Smirnov and F. Vissani, *Nucl. Phys.* **B460** (1996) 37; D. Du and C. Liu, *Mod. Phys. Lett.* **A10** (1995) 1837; C. Liu, *Mod. Phys. Lett.* **A12** (1997) 329; S. Roy and B. Mukhopadhyaya, *Phys. Rev.* **D55** (1997) 7020; B. Mukhopadhyaya, S. Roy and F. Vissani, *Phys. Lett.* **B443** (1998) 191; G. Dvali and Y. Nir, *JHEP* **9810:014**,1998 ; M. Bisset, O. C. W. Kong, C. Macesanu, L. H. Orr, *Phys. Rev. D* **62** (2000) 035001.
- [10] M. Hirsch, M. A. Diaz, W. Porod, J. C. Romao and J. W. F. Valle, *Phys. Rev. D* **62** (2000) 113008. Also see, by the same authors, *Phys. Rev.***D61** (2000) 071703; hep-ph/0009066; hep-ph/0009127.
- [11] D. E. Kaplan and A. E. Nelson, *JHEP* **033:0001**, 2000.
- [12] R. Kitano and K. Oda, *Phys. Rev D* **61** (2000) 113001.
- [13] S. Rakshit, G. Bhattacharyya and A. Raychaudhuri, *Phys. Rev.* **D59** (1999) 091701. A. Abada

- and M. Losada, *Nucl. Phys.* **B 585** (2000) 45; R. Adhikari and G. Omanovic, hep-ph/9802390; *Phys. Rev.* **D 59** (1999) 073003; O. Haug, J. D. Vergados, A. Faessler and S. Kovalenko, *Nucl. Phys.* **B 565** (2000) 38; R. Adhikari, A. Sil and A. Raychaudhuri, hep-ph/0105119.
- [14] M. Drees, S. Pakavasa, X. Tata and T. ter Veldhuis, *Phys. Rev.* **D 57** (1998) R5335.
- [15] A. S. Joshipura and S. K. Vempati, *Phys. Rev.* **D60** (1999) 111303.
- [16] B. de Carlos and P. L. White, *Phys. Rev.* **D54** (1996) 3427; E. Nardi, *Phys. Rev.* **D55** (1997) 5772.
- [17] G. Gamberini, G. Ridolfi and F. Zwirner, *Nucl. Phys.* **B331** (1990) 331.
- [18] E. J. Chun, S. K. Kang, *Phys. Rev.* **D61**(2000) 075012; E. J. Chun, hep-ph/0001236.
- [19] F. Borzumati, hep-ph/9702307.
- [20] A. S. Joshipura and M. Nowakowski, *Phys. Rev.* **D51** (1995) 2421.
- [21] M. Dine, A. E. Nelson and Y. Shirman, *Phys. Rev.* **D 51** (1995) 1362; M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, *Phys. Rev.* **D 53** (1996) 2658. For original references and reviews please see, G. F. Giudice and R. Rattazzi, *Phys. Rep.* **322** (1999) 419; S. L. Dubovsky, D. S. Gorbunov and S. V. Troitsky, *Phys. Usp.* **42** (1999) 623, [hep-ph/9905466].
- [22] K. Choi, K. Hwang and E. J. Chun, *Phys. Rev.* **D 60** (1999) 031301 and Ref. [18].
- [23] M. Dine, Y. Nir and Y. Shirman, *Phys. Rev.* **D 55** (1997) 1501; K. S. Babu, C. Kolda and F. Wilczek, *Phys. Rev. Lett.* **77** (1996) 3070; S. Dimopoulos, S. Thomas and J. D. Wells, *Nucl. Phys.* **B488** (1997) 39; J. A. Bagger, K. Matchev, D. M. Pierce and R. Zhang, *Phys. Rev.* **D55** (1997) 3188.
- [24] S. P. Martin , *Phys. Rev.* **D 55** (1997) 3177
- [25] U. Sarid and R. Rattazzi, *Nucl. Phys.* **B 501** (1997) 297.
- [26] A. S. Joshipura, V. Ravindran and S. K. Vempati, *Phys. Lett.* **B451** (1999) 98.
- [27] R. Barate *et. al*, [ALEPH Collaboration], *Eur. Phys. J* **C 16** (2000) 71.

- [28] M. C. Gonzalez-Garcia, M. Maltoni, C. Peña-Garay and J. W. F. Valle, *Phys. Rev. D* **63** (2001)(033005); J. Bahcall, P. I. Krastev and A. Yu. Smirnov, *JHEP* **0105:015**(2001).
- [29] Y. Oyama *et. al.*, [Super-Kamiokande Collaboration], hep-ex/0104015.
- [30] Two generation analyses including the recent results from SNO (nucl-ex/0106015) have been presented in, G. L. Fogli *et. al.*, hep-ph/0106247; J. N. Bahcall *et. al.*, hep-ph/0106258; A. Bandyopadhyay *et. al.*, hep-ph/0106264. Also see, V. Barger *et. al.*, hep-ph/0106207.
- [31] N. K. Falck, *Zeit. fur Physik* **C30** (1986) 247.
- [32] For a list of MSSM RGE, please see, L. E. Ibanez and C. Lopez, *Nucl. Phys.* **B233** (1984) 511; M. Carena *et. al.* *Nucl. Phys.* **B 491** (1997) 103.
- [33] A. S. Joshipura, R. D. Vaidya and S. K. Vempati, *Phys. Rev D* **62** (2000) 093020.






FIG. 1. Neutrino mass eigen value ratio  $\frac{m_{\nu_2}}{m_{n_3}}$  plotted versus  $\Lambda$  (GeV) assuming trilinear  $\lambda'$  couplings of similar strength to be the only source of  $R$  violation.  $\Lambda$  is defined in the text.




FIG. 2. The neutrino mass ratio  $\frac{m_{loop}}{m_0}$  plotted versus  $\Lambda$  (GeV) assuming trilinear  $\lambda$  couplings of similar strength to be the only source of  $R$  violation.